



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON A COLLECTION OF BIRDS MADE ON THE
ISLANDS OF ANGUILLA AND CAY SAL OR SALT
CAY, BAHAMA ISLANDS, BY MR. CYRUS
S. WINCH, DURING MAY, 1891.

BY CHARLES B. CORY.

THE Islands of Anguilla and Cay Sal are situated about sixty miles north of Cuba at the extreme southwestern portion of the Bahama Banks. Anguilla is high and rather barren, the lower portions being covered with a low scrub and brush. Unfortunately much of the vegetation had been destroyed by fire, and portions of it were still burning at the time of Mr. Winch's visit. He observed many birds passing on their migration north. On May 10 and 11 the Bobolink and Black-poll Warbler (*Dendroica striata*) were abundant. The species taken on Anguilla were:

<i>Sterna fuliginosa</i> Gmel.	<i>Chelidon erythrogaster</i> (Bodd.).
<i>Zenaida zenaida</i> (Bonap.).	<i>Setophaga ruticilla</i> (Linn.).
<i>Coccyzus americanus</i> (Linn.).	<i>Geothlypis trichas</i> (Linn.).
<i>Dolichonyx oryzivorus</i> (Linn.).	<i>Dendroica cærulescens</i> (Gmel.).
<i>Callichelidon cyaneoviridis</i> Bryant.	<i>Dendroica striata</i> (Forst.).

Cay Sal is rather more fertile than Anguilla and has a fresh water pond in the interior surrounded by cocoanut and other trees. A large rabbit (probably introduced) was at one time abundant, but is claimed now to be dying off. The Burrowing Owl on this island appears to be the same as the Florida form. Mr. Winch reached Cay Sal on May 14 and found birds abundant, mostly our common species on their migration. On May 16 birds were less numerous and on May 19 very few were to be seen on the island. He procured the following:

<i>Sterna antillarum</i> (Less.).	<i>Callichelidon cyaneoviridis</i> Bryant.
<i>Arenaria interpres</i> (Linn.).	<i>Seiurus noveboracensis</i> (Gmel.).
<i>Speotyto cunicularia floridana</i> Ridgw.—Resident and not uncommon.	<i>Geothlypis trichas</i> (Linn.).
<i>Coccyzus americanus</i> (Linn.).	<i>Geothlypis agilis</i> (Wils.).—Not previously recorded from the West Indies.
<i>Tyrannus dominicensis</i> (Gmel.).	<i>Dendroica vigorsii</i> (Aud.).
<i>Agelaius phæniceus bryanti</i> Ridg.	<i>Turdus aliciae bicknelli</i> (Ridgw.).
<i>Dolichonyx oryzivorus</i> (Linn.).	—First record for the West Indies.
<i>Chelidon erythrogaster</i> (Bodd.).	